
مداخلات ترک سیگار برای مسلولین

راهنمای سال ۲۰۰۸ برای کشورهای کم درآمد

مرکز تحقیقات پیشگیری و کنترل دخانیات
پژوهشکده سل و بیماریهای ریوی
دانشگاه علوم پزشکی شهید بهشتی
جمعیت مبارزه با استعمال دخانیات ایران

مترجم:

پریزاد سینایی

ویراستار:

دکتر هومن شریفی

دکتر محمد ورهرام

زیر نظر:

دکتر محمدرضا مسجدی

Slama, Karen

978-600-5069-13-6

RC /

Tobacco Cessation Interventions for Tuberculosis Patients

:

5

8

:

15

18

18

19

:

:

5Rs 5As

:

:

১৭

:

১৮

:

Mira Aghi, Natasha Herrera, Jacques Prignot

International Journal of Tuberculosis and Lung Disease

)

(

:

PPD-S 5-TU (\geq mm)

(/)

mm

cohort

:

)
(

cohort

Tamil Nadu nested

BCG

« »

/) (/)
((/)

Mycobacterium vaccae

Cox

MDR-TB

cohort

acid-fast

(%)

cohort

/

Catalonia

Recife

MDR-TB

MDR-TB

MDR-TB

Isoniazid

Nicotiana

:

(/)

DSM-IV-TR

/

)

:

(

« »

;

%

%

« »

« »

« »

(: (

()

:

(
(

(
(
(
(
(
(
(

:

(
/
(
(

:

:

:

:

()

:

.

:

.

.

.

.

.

.

.

.

.

.

.

.

«

»

:

-
-
-
-
-
-
-
-

« »

»

«

:

)

(

()

()

(

(

(National Tuberculosis Program)

:

:

.

.

.

.

.

.

.

:

:

.

⋮
»

«

« »

/

()

« » « »

()

:5Rs 5As

: 5A

:Ask

:Advise

5Rs

:Asses

:Assist

:Arrange

5Rs

:Relevance

:Risks

:Rewards

:Roadblocks

)
(.

:Repetition

(Fagerström)

(/)

: /

*

≥

≤

≥

≥

*

:

() .

:

()

:

;

)
() .(

()

;

:

:

(

(

(

(

:

/

()

(

(

(

؛
:
(
) (
(
(
(

: /

:

(/)

•

•

•

: /

~

*

= = = = = *

:

)

(

:

() .

-
-
-
-

-
-
-
-
-

-
-
-
-
-

:

:

/

:

: /

:

:

)

(

:

:

:

:

(

(

(

cohort

؛
:

-
-
-
-
-
-
-

؛

(:)

-
-

:
:
()
:
()
()
:
()
:
()

« »

:

-
-
-
-
-

:

()

:

- (
- (
- (

(

(

(%)

:

:

•

:

•

•

« »

< ()

•

()

:

•

* ()

:

•

()

:

•

" "

" "

Fagerström

()

:

•

•

•

:

•

*

.

()

.

: •

.

.

()

: •

: •

:

: •

.()

:

: •

.

:

: •

()

()

: •

)

:

: •

(

:

: •

Fagerström

:

() .

;
()

- : •
- : •
- : •

: •

" "

: •

: •

: •

" " " "

;
()
" "

(.

	:	:	●
" "	" "	-	
" " " "	" "	-	
	:	:	●
	:	<	
	:	:	●
	:	:	●
	:	:	●
" " " "	:	:	●
" " " "	:	:	●
	:	:	●

- : :
.
- : :
.
- : :
.
- : :
.

:

:

Abal AT, Jayakrishnana B, Parwer S, El Shamy A, Abahussain E, Sharma PN. Effect of cigarette smoking on sputum smear conversion in adults with active pulmonary tuberculosis. *Respir Med* 2005; 99: 415–420.

Alcaide J, Altet MN, Plans P, Parron I, Foguera L, Salto E, Dominguez A, Pardell H, Salleras L. Cigarette smoking as a risk factor for tuberculosis in young adults: a case-control study. *Tubercle Lung Dis* 1996; 77: 112–116.

Altet-Gomez MN, Alcaide J, Godoy P, Romero MA, Hernandez del Rey I. Clinical and epidemiological aspects of smoking and tuberculosis: a study of 13038 cases. *Int J Tuberc Lung Dis* 2005; 9: 430–436.

Altet MN, Alcaide J, Plans P, Taberner JL, Salto E, Folguera LI, Salleras L. Passive smoking and risk of pulmonary tuberculosis in children immediately following infection. A case-control study. *Tubercle Lung Dis* 1996; 77: 537–544.

Anderson RH, Sy FS, Thompson S, Addy C. Cigarette smoking and tuberculin skin test conversion among incarcerated adults. *Am J Prev Med* 1997; 13: 175–181.

Barroso EC, Mota RMS, Santos RO, Sousa ALO, Barroso JB, Rodrigues JLN. Risk factors

for acquired multidrug-resistant tuberculosis. *J Pneumol* 2003; 29: 89–97.

Brown KE, Campbell AH. Tobacco, alcohol and tuberculosis. *Brit J Dis Chest* 1961; 55: 150–158.

Buskin SE, Gale JL, Weiss NS, Nolan CM. Tuberculosis risk factors in adults in King County, Washington, 1988–1990. *Am J Public Health* 1994; 84: 1750–1756.

Chiang C-Y, Chang C-T, Chang R-E, Li C-T, Huang R-M. Patient and health system delays in the diagnosis and treatment of tuberculosis in Southern Taiwan. *Int J Tuberc Lung Dis* 2005; 9: 1006–1012.

Chang KC, Leung CC, Tam CM. Tuberculosis risk factors in a silicotic cohort in Hong Kong. *Int J Tuberc Lung Dis* 2001; 5: 177–184.

den Boon S, van Lill, SWP, Borgdorff MW, Verver S, Batemen ED, Lombard CJ, Enarson DA, Beyers N. The association between smoking and tuberculosis infection: a population survey in a high tuberculosis incidence area. *Thorax* 2005; 60: 555–557.

Doll R, Hill AB. The mortality of doctors in relation to their smoking habits. *BMJ* 1954; reprinted in *BMJ* 2002; 328: 1529–1533.

Doll R, Peto R, Wheatley K, Gray R, Sutherland I. Mortality in relation to smoking: 40 years' observations on male British doctors. *BMJ* 1994; 309: 901–911.

Dos Santos MA, Albuquerque MF, Ximenes RA, Lucena-Silva NL, Braga C, Campelo AR, Dantas OM, Montarroyos UR, Souza WV, Kawasaki AM, Rodriguez LC.

Risk factors for treatment delay in pulmonary tuberculosis in Recife, Brazil.

BMC Public Health 2005; 5: 25.

Durban Immunotherapy Trial Group. Immunotherapy with *Mycobacterium vaccae* in patients with newly diagnosed pulmonary tuberculosis: a randomised controlled trial. *Lancet* 1999; 354: 116–119.

52 REFERENCES

- Gajalakshmi V, Peto R, Kanaka TS, Jha R. Smoking and mortality from tuberculosis and other diseases in India: retrospective study of 43,000 adult male deaths and 35,000 controls. *Lancet* 2003; 362: 507–515.
- Hnizdo E, Murray J. Risk of pulmonary tuberculosis relative to silicosis and exposure to silica dust in South African gold miners. *Occup Environ Med* 1998; 55: 496–502.
- Hussain H, Akhtar S, Nanan D. Prevalence of and risk factors associated with *Mycobacterium tuberculosis* infection in prisoners, North West Frontier Province, Pakistan. *Int J Epidemiol* 2003; 32: 794–799.
- Kolappan C, Gopi PG. Tobacco smoking and pulmonary tuberculosis. *Thorax* 2002; 57: 964–966.
- Kuemmerer JM, Comstock GW. Sociologic concomitants of tuberculin sensitivity. *Am Rev Respir Dis* 1967; 96: 885–892.
- Lam TH, Ho SY, Hedley AJ, Mak KH, Peto R. Mortality and smoking in Hong Kong: case control study of all adult deaths in 1998. *BMJ* 2001; 323: 1–6.
- Lewis JG, Chamberlain DA. Alcohol consumption and smoking habits in male patients with pulmonary tuberculosis. *Brit J Prev Soc Med* 1963; 17: 149–152.
- Leung CC, Yew WW, Chan CK, Tam CM, Lam CW, Chang KC, Chau CH, Lau KS, Law WS. Smoking and tuberculosis in Hong Kong. *Int J Tuberc Lung Dis* 2003; 7: 980–986.
- Leung CC, Li T, Lam TH, Yew WW, Law WS, Tam CM, Chan WM, Chan CK, Ho KS, Chang KC. Smoking and tuberculosis among the elderly in Hong Kong. *Am J Respir Crit Care Med* 2004; 170: 1027–1033.
- Lienhardt C, Fielding K, Sillah JS, Bah B, Gustafson P, Warndorff D, Palayew M, Lisse I, Donkor S, Diallo S, Manneh K, Adegbola R, Aaby P, Bah-Sow O, Bennett S, McAdam K. Investigation of the risk factors for tuberculosis: a case-control study in three countries in West Africa. *Int J Epidemiol* 2005; 34: 914–923.
- Liu B-Q, Peto R, Chen Z-M, Boreham J, Wu Y-P, Li J-Y, Campbell TC, Chen J-S. Emerging tobacco hazards in China: 1. Retrospective proportional mortality study of one million deaths. *BMJ* 1998; 317: 1411–1422.
- Lowe CR. An association between smoking and respiratory tuberculosis. *BMJ* 1956; 2: 1081–1086. Same data further analysed in Edwards JH. Contribution of cigarette smoking to respiratory disease. *Br J Prev Soc Med* 1957; 11: 10–21.
- Nisar M, Williams CS, Ashby D, Davies G. Tuberculin testing in residential homes for the elderly. *Thorax* 1993; 48: 1257–1260.
- Perez-Padilla R, Perez-Guzman C, Baez-Saldana R, Torres-Cruz A. Cooking with biomass stoves and tuberculosis: a case-control study. *Int J Tuberc Lung Dis* 2001; 5: 441–447.
- Plant AJ, Watkins RE, Gushulak B, O'Rourke T, Jones W, Streeton J, Sang D. Predictors of tuberculin reactivity among prospective Vietnamese migrants: the effect

of smoking. *Epidemiol Infect* 2002; 128: 37–45.

Ruddy M, Balabanova Y, Graham C, Fedorin I, Malomanova N, Elisarova E, Kuznetznov S, Gusarova G, Zakharova S, Melentyev A, Krukova E, Golishevskaya V, Erokhin V, Dorozhkova I, Drobniowski F. Rates of drug resistance and risk factor analysis in civilian and prison patients with tuberculosis in Samara Region, Russia. *Thorax* 2005; 60: 130–135.

Santha T, Garg R, Frieden TR, Chandrasekaran V, Subramani R, Gopi PG, Selvakumar N, Ganapathy S, Charles N, Rajamma J, Narayanan PR. Risk factors associated with default, failure and death among tuberculosis patients treated in a

REFERENCES 53

DOTS programme in Tiruvallur district, South India, 2000. *Int J Tuberc Lung Dis* 2002; 6: 780–788.

Sitas F, Urban M, Bradshaw D, Kielkowski D, Bah S, Peto R. Tobacco attributable deaths in South Africa. *Tob Control* 2004; 13: 396–399.

Singh M, Mynak ML, Kumar L, Mathew JL, Jindal SK. Prevalence and risk factors for transmission of infection among children in household contact with adults having pulmonary tuberculosis. *Arch Dis Child* 2005; 90: 624–628.

Slama K, Chiang CY, Enarson DA, Hassmiller K, Fanning A, Gupta PC, Ray CS. Tobacco

and tuberculosis: a qualitative systematic review and meta-analysis. *Int J Tuberc Lung Dis* 2007; 11: 1049–1061.

Tekkel M, Rahu M, Loit H-M, Baburin A. Risk factors for pulmonary tuberculosis in Estonia. *Int J Tuberc Lung Dis* 2002; 6: 887–894.

Thomas A, Gopi PG, Santha T, Chandrasekaran V, Subramani R, Selvkumar N, Eusuff SI, Sadacharam K, Narayanan PR. Predictors of relapse among pulmonary tuberculosis patients treated in a DOTS programme in South India. *Int J Tuberc Lung Dis* 2005; 9: 556–561.

Tipayamongkhogul M, Podhipak A, Chearskul S, Sunakorn P. Factors associated with the development of tuberculosis in BCG immunized children. *Southeast Asian J Trop Med Public Health* 2005; 36: 145–150.

Tocque K, Bellis MA, Beeching JN, Syed Q, Remington T, Davies PDO. A casecontrol

study of lifestyle risk factors associated with tuberculosis in Liverpool, North-West England. *Eur Respir J* 2001; 18: 959–964.

Watkins RE, Plant AJ. Does smoking explain sex differences in the global tuberculosis epidemic? *Epidemiol Infect* 2006; 134: 333–339.

Webb GB. The effect of the inhalation of cigarette smoke on the lungs. A clinical study. *Am Rev Tuberc* March 1918; 25–27.

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*.

Fourth Edition. Text revision. Washington, DC, USA: American Psychiatric Association, 2000.

- Boyle P, Gandini S, Robertson C, Zatonski W, Fagerstrm K, Slama K, Kunze M, Gray N. Characteristics of smokers attitudes towards stopping : survey of 10,295 smokers in representative samples from 17 European cities. *Eur J Public Health* 2000; 10: 5–14.
- Callard C, Thompson D, Collishaw N. Curing the addiction to profits: a supplyside approach to phasing out tobacco. Ottawa, ON, Canada: Canadian Centre for Policy Alternatives (CCPA), 2005.
- Dalton MA, Sargent JD, Beach ML, Titus-Ernstoff L, Gibson JJ, Ahrens MB, Tickle JJ, Heatherton TF. Effect of viewing smoking in movies on adolescent smoking initiation: a cohort study. *Lancet* 2003; 362: 281–285.
- Difranza JR, Rigotti NA, McNeill AD, Ockene JK, Savageau JA, St Cyr D, Coleman M. Initial symptoms of nicotine dependence in adolescents. *Tob Control* 2000; 9: 313–319.
- Difranza JR, Savageau JA, Rigotti NA, Fletcher K, Ockene JK, McNeill AD, Coleman M, Wood C. Development of symptoms of tobacco dependence in youths: 30 month follow up data from the DANDY study. *Tob Control* 2002; 11: 228–235.
- 54 REFERENCES**
- Gervais A, O’Loughlin J, Meshefedjian G, Bancenj C, Trembley M. Milestones in the natural course of onset of cigarette use among adolescents. *CMAJ* 2006; 175: 255–261.
- Goodman J. Tobacco in history. The cultures of dependence. New York, NY, USA: Routledge, 1994.
- Guindon GE, Boisclair D. Past, current and future trends in tobacco use. *Health Nutrition and Population Discussion Paper, Economics of tobacco control. Paper no. 6.* Washington, DC, USA: The World Bank, 2003.
- Hughes JR, Higgins ST, Bickel WK. Nicotine withdrawal versus other drug withdrawal syndromes: similarities and dissimilarities. *Addiction* 1994; 89: 1461–1470.
- Hughes JR, Keely J, Naud S. Shape of the relapse curve and long-term abstinence among untreated smokers. *Addiction* 2004; 99: 29–38.
- Hyland A, Borland R, Li Q, Yong HH, McNeill A, Fong GT, O’Connor RJ, Cummings KM. Individual-level predictors of cessation behaviours among participants in the International Tobacco Control (ITC) Four Country Survey. *Tob Control* 2006; 15 (Suppl 3): iii83–94.
- International Agency for Research on Cancer. Tobacco smoke and Involuntary smoking. Vol 83. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Lyon, France: IARC, 2004: pp 1191–1203.
- LeHouezec J. Role of nicotine pharmacokinetics in nicotine addiction and nicotine replacement therapy: a review. *Int J Tuberc Lung Dis* 2003; 7: 811–819.
- Mowery PD, Farrelly MC, Haviland ML, Gable JM, Wells HE. Progression to established smoking among US youths. *Am J Public Health* 2004; 94: 331–337.
- Prochaska JO. Moving beyond the transtheoretical model. *Addiction* 2006; 101: 768–774.
- Rose JE, Behm FM, Levin ED. Role of nicotine dose and sensory cues in the regulation of smoke intake. *Pharmacol Biochem Behav* 1993; 44: 891–900.

- Royal College of Physicians. Nicotine addiction in Britain: a report of the Tobacco Advisory Group of the Royal College of Physicians. London, UK: Royal College of Physicians, 2000.
- Slama K. Current challenges in tobacco control. *Int J Tuberc Lung Dis* 2004; 10: 1160–1172.
- Slama K. Tobacco control and prevention. A guide for low-income countries. Paris, France: International Union Against Tuberculosis and Lung Disease, 1998.
- Tucker JS, Ellickson PL, Klein DJ. Smoking cessation during the transition from adolescence to young adulthood. *Nicotine Tob Res* 2002; 4: 321–332.
- US Department of Health and Human Services. The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General. Atlanta, GA, USA: US Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2006.
- World Health Organization. Biobehavioural processes underlying dependence. In: *Neuroscience of psychoactive substance use and dependence*. Geneva, Switzerland: WHO, 2004: pp 43–65.
- da Costa e Silva V, ed. Policy recommendations for smoking cessation and treatment of tobacco dependence. (Advancing tobacco control in the XXIst century). Geneva, Switzerland: World Health Organization, 2003: pp 1–10.
- REFERENCES 55**
- World Health Organization. Psychopharmacology of dependence for different drug classes. In: *Neuroscience of psychoactive substance use and dependence*. Geneva, Switzerland: WHO, 2004: pp 75–78.
- World Health Organization. WHO Framework Convention on Tobacco Control. Geneva, Switzerland: WHO, 2003. <http://www.who.int/tobacco/framework/en/>

- Auer C, Sarol J Jr, Tanner M, Weiss M. Health seeking and perceived causes of tuberculosis among patients in Manila, Philippines. *Trop Med Int Health* 2000; 5: 648–656.
- Bass F. Mobilizing physicians to conduct clinical intervention in tobacco use through a medical-association program: 5 years' experience in British Columbia. *Can Med Assoc J* 1996; 154: 159–163.
- Carlos RC, Fendrick AM. Improving cancer screening adherence: using the “teachable moment” as a delivery setting for educational interventions. *Am J Manag Care* 2004; 10: 247–248.
- Chiang CY, Slama K, Enarson D. Associations between tobacco and tuberculosis. *Int J Tuberc Lung Dis* 2007; 11: 258–262.
- Cohen SJ, Halvorson HW, Gosselink CA. Changing physician behavior to improve disease prevention. *Prev Med* 1994; 23: 284–291.

da Costa e Silva V, ed. Policy recommendations for smoking cessation and treatment of tobacco dependence (Advancing tobacco control in the XXIst century). Geneva, Switzerland: World Health Organization, 2003.

Elford RW, Yeo M, Jennett PA, Sawa RJ. A practical approach to lifestyle change counselling in primary care. *Patient Educ Couns* 1994; 24: 175–183.

El Sony A, Slama K, Salieh M, Elhaj H, Adam K, Hassan A, Enarson DA. Feasibility of brief tobacco cessation advice for tuberculosis patients. A study from Sudan. *Int J Tuberc Lung Dis* 2007; 11: 150–155.

Lancaster T, Stead L. Physician advice for smoking cessation. *Cochrane Database Syst Rev* 2004; (4): CD000165.

Shiffman S. Reflections on smoking relapse research. *Drug Alc Rev* 2006; 25: 15–20.

Slama K. Behavioural approaches to cessation. In: Lu R, Mackay J, Niu S, Peto R, eds. *Tobacco: the growing epidemic*. London, UK: Springer-Verlag, 2000.

De Leon J, Diaz FJ, Becona E, Gurpegui M, Jurado D, Gonzalez-Pinto A. Exploring brief measures of nicotine dependence for epidemiological surveys. *Addict Behav* 2003; 28: 1481–1486.

El Sony A, Slama K, Salieh M, Elhaj H, Adam K, Hassan A, Enarson DA. Feasibility of brief tobacco cessation advice for tuberculosis patients. A study from Sudan. *Int J Tuberc Lung Dis* 2007; 10: 150–155.

Fiore MC, Bailey WC, Cohen SJ, et al. Treating tobacco use and dependence. Clinical practice guideline. Rockville, MD, USA: US Department of Health and Human Services, Public Health Service, June 2000.

Foulds J, Steinberg MB, Williams JM, Ziedonis DM. Developments in pharmacotherapy for tobacco dependence: past, present and future. *Drug Alcohol Rev* 2006; 25: 59–71.

56 REFERENCES

Lando H, Hennrikus D, McCarty M, Vessey J. Predictors of quitting in hospitalized smokers. *Nicotine Tob Res* 2003; 5: 215–222.

Le Houezec J. Role of nicotine pharmacokinetics in nicotine addiction and nicotine replacement therapy: a review. *Int J Tuberc Lung Dis* 2003; 7: 811–819.

Okuyemi KS, Nollen NL, Ahluwalia JS. Interventions to facilitate smoking cessation. *Am Acad Family Physicians* 2006; 74: 262–271.

Piper ME, McCarthy DE, Baker TB. Assessing tobacco dependence: a guide to measure evaluation and selection. *Nicotine Tob Res* 2006; 8: 339–351.

Roddy E, Antoniak M, Britton J, Molyneux A, Lewis S. Barriers and motivators to accessing smoking cessation services amongst deprived smokers—a qualitative study. *BMC Health Serv Res* 2006; 6: 147.

Slama K. Behavioural approaches to cessation. In: Lu R, Mackay J, Niu S, Peto R, eds. *Tobacco: the growing epidemic*. London, UK: Springer-Verlag, 2000.

Solberg LI, Maciosek MV, Edwards NM, Khanchandani HS, Goodman MJ. Repeated tobacco-use screening and intervention in clinical practice: health impact and cost effectiveness. *Am J Prev Med* 2006; 31: 62–71.

US Department of Health and Human Services. Reducing tobacco use: a report of

the Surgeon General. Atlanta, GA, USA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2000.

Wellman R, Savageau J, Godiwala S, Savageau N, Friedman K, Hazelton J, DiFranza J. A comparison of the Hooked on Nicotine Checklist and the Fagerstrm Test for Nicotine Dependence in adult smokers. *Nicotine Tob Res* 2006; 8: 575–580.

World Health Organization. Biobehavioural processes underlying dependence. In: *Neuroscience of psychoactive substance use and dependence*. Geneva, Switzerland: WHO, 2004: pp 43–66.

American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*.

4th ed. Text revision. Washington, DC, USA: APA, 2000.

Axelrod S. Smoking cessation through functional analysis. *J Applied Behav Anal* 1991; 24: 717–718.

Carroll KM. *Cognitive-behavioral therapy for cocaine users. Manual 1*. NIH Publication Number 98-4308. Bethesda, MD, USA: National Institute on Drug Abuse, 1998.

Cummings KM, Giovino G, Jaen CR, Emrich LJ. Reports of smoking withdrawal symptoms over a 21 day period of abstinence. *Addict Behav* 1985; 10: 373–381.

Fiore MC, Bailey WC, Cohen SJ, Dorfman SF, Goldstein MG, Gritz ER, et al. *Treating tobacco use and dependence. Clinical practice guideline*. Rockville, MD, USA: US Department of Health and Human Services. Public Health Service. June 2000.

Hogarth L, Dickinson A, Hutton SB, Bamborough H, Duka T. Contingency knowledge is necessary for learned motivated behaviour in humans: relevance for addicted behaviour. *Addiction* 2006; 101: 1153–1166.

Hughes JR. Clinical significance of tobacco withdrawal. *Nicotine Tob Res* 2006; 8: 153–156.

REFERENCES 57

Kadowaki T, Watanabe M, Okayama A, Hishida K, Okamura T, Miyamatsu N, Hayakawa

T, Kita Y, Ueshima H. Continuation of smoking cessation and following weight change after intervention in a healthy population with high smoking prevalence. *J Occup Health* 2006; 48: 402–406.

Myers MG, Brown SA, Kelly JF. A cigarette smoking intervention for substanceabusing adolescents. *Cogn Behav Pract* 2000; 7: 64–82.

National Institute on Drug Abuse. *Principles of drug addiction treatment: a researchbased guide*. NIH Publication No. 99-4180. Bethesda, MD, USA: NIDA, 1998: pp 15.

Park E-W, Tudiver F, Schultz JK, Campbell T. Does enhancing partner support and

- interaction improve smoking cessation? A meta-analysis. *Ann Fam Med* 2004; 2: 170–174.
- Pednekar MS, Gupta PC, Shukla HC, Hebert JR. Association between tobacco use and body mass index in urban Indian population: implications for public health in India. *BMC Public Health* 2006; 6: 70.
- Shiffman S. Reflections on smoking relapse research. *Drug Alc Rev* 2006; 25: 15–20.
- Slama K. Health behaviour and change. In: Sancho-Garnier H, Anderson A, Biedermann A, Lynge E, Slama K, eds. Evidence-based cancer prevention: strategies for NGOs. A UICC Handbook for Europe. Geneva, Switzerland: International Union Against Cancer, 2004: pp 48–59.
- Slama K, Chiang CY, Enarson DA. Tobacco cessation and brief advice. *Int J Tuberc Lung Dis* 2007; 11: 612–616.
- US Department of Health and Human Services. Reducing tobacco use: a report of the Surgeon General. Atlanta, GA, USA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2000.
- World Health Organization. Psychopharmacology of dependence for different drug classes. In: Neuroscience of psychoactive substance use and dependence. Geneva, Switzerland: WHO, 2004: pp 75–78.

